

Localization with *Less Larmes*: Simply MSA

Victor Chulaevsky¹

Département de Mathématiques et d'Informatique,
Université de Reims, Moulin de la Housse, B.P. 1039,
51687 Reims Cedex 2, France
E-mail: victor.tchoulaevski@univ-reims.fr

December 2008

Abstract: We give a short summary of the fixed-energy Multi-Scale Analysis (MSA) of the Anderson tight binding model in dimension $d \geq 1$ and show that this technique admits a straightforward extension to multi-particle systems. We hope that this short note may serve as an elementary introduction to the MSA.

1. Introduction

In this paper, we study spectral properties of random lattice Schrödinger operators at a fixed, but arbitrary, energy $E \in \mathbb{R}$, in the framework of the MSA. The idea of the fixed-energy scale induction goes back to [FS83], [Dr87]¹ and [S88]. While the fixed-energy analysis alone does not allow to prove spectral localization, it provides a valuable information. Besides, from the physical point of view, a sufficiently rapid decay, with probability one, of Green functions in finite volumes, combined with the celebrated Kubo formula for the zero-frequency conductivity $\sigma(E)$, shows that $\sigma(E) = 0$ for the disordered systems in question.

The main motivation for studying in this paper only the fixed-energy properties of random Hamiltonians came from an observation that such analysis can be made very elementary, even for multi-particle systems considered as difficult since quite a long time (cf. recent works [CS08], [CS09A], [AW08], [CS09B], [BCSS08], [BCS08]).

An earlier version of this manuscript represented an extended variant of the talk given by the author of these lines in the framework of the program MPA run by the Isaac Newton Institute in 2008. Questions and remarks made by participants to the program made it clear that the brevity of the first version was sometimes excessive, and that additional illustrations and explanations would be useful. Such modifications are partially implemented in the present version, and

¹ I thank A. Klein and T. Spencer for pointing out the PhD thesis by H. von Dreifus [Dr87] (scientific advisor T. Spencer) where the fixed-energy approach had been used.

it will probably evolve further in order to make this (very) short introduction to the MSA fairly clear not only for mathematicians, but also for researchers with different backgrounds, interested more by results and basic ideas than by formal constructions. Producing clear illustrations is a time-consuming process, but live discussions evidence that they can be more useful than equations.

2. The models and some geometric definitions

A popular form of a single-particle Hamiltonian in presence of a random external potential $gV(x; \omega)$ is as follows:

$$H(\omega) = -\Delta + gV(x; \omega), \quad (2.1)$$

where the parameter $g \in \mathbb{R}$ is often called the coupling constant and Δ is the nearest-neighbor lattice Laplacian:

$$(\Delta f)(x) = \sum_{y: \|y-x\|=1} f(y), \quad x, y \in \mathbb{Z}^d,$$

and $V(x; \omega)$ acts as a multiplication operator on $\mathbb{Z}^d$. For the sake of simplicity, the random field $\{V(x; \omega), x \in \mathbb{Z}^d\}$ will be assumed IID, although a large class of correlated random fields can also be considered. In this paper, we consider only the case of "large disorder", i.e., we assume that $|g|$ is sufficiently large, although the case of "low energies" can also be considered. An IID random field on $\mathbb{Z}^d$ is completely determined by its marginal probability distribution at any site $x \in \mathbb{Z}^d$, e.g., for $x = 0$. We assume that the marginal distribution function of potential V (defined by $F_V(t) = \mathbb{P}\{V(0; \omega) \leq t\}$, $t \in \mathbb{R}$) is Hölder-continuous:

$$\forall t \in \mathbb{R}, \forall \epsilon \in (0, 1] \quad F_V(t + \epsilon) - F_V(t) \leq \text{Const} \epsilon^b, \quad (2.2)$$

for some $b > 0$.

For $N > 1$ particles, positions of which will be denoted by $x_1, \dots, x_N$, or, in vector notations, $\mathbf{x} = (x_1, \dots, x_N) \in \mathbb{Z}^{Nd}$, we introduce an interaction energy $U(\mathbf{x})$. Again, for the sake of simplicity of presentation, we assume that

$$U(\mathbf{x}) = U(x_1, \dots, x_N) = \sum_{1 \leq j < k \leq N} U_2(\|x_j - x_k\|), \quad (2.3)$$

where $U_2(r)$, $r \geq 0$, is a bounded two-body interaction potential. The N -particle Hamiltonian considered below will have the following form:

$$H^{(N)}(\omega) = \sum_{j=1}^N \left(-\Delta^{(j)} + gV(x_j; \omega) \right) + U(\mathbf{x}). \quad (2.4)$$

Given a lattice subset $\Lambda \subset \mathbb{Z}^d$, we will work with subsets thereof called boxes. It is convenient to allow boxes $\Lambda_\ell(u) \subset \Lambda$ of the following form:

$$\Lambda_\ell(u) = \{x : \|x - u\| \leq \ell - 1\},$$

where $\|\cdot\|$ is the sup-norm: $\|x\| = \max_{1 \leq j \leq d} |x_j|$. Further, we introduce a notion of internal and external "boundaries" relative to Λ :

$$\begin{aligned}\partial^- \Lambda_\ell(u) &= \{x \in \Lambda : \|x - u\| \leq \ell - 1\} \\ \partial^+ \Lambda_\ell(u) &= \{x \in \Lambda : \text{dist}(x, \Lambda_\ell(u)) = 1\}.\end{aligned}$$

We also define the boundary $\partial \Lambda_\ell(u) \in \Lambda$ by

$$\partial \Lambda_\ell(u) = \{(x, x') : x \in \partial^- \Lambda_\ell(u), x' \in \partial^+ \Lambda_\ell(u), \|x - x'\| = 1\}.$$

By resolvent identity, if operators A and $A + B$ are invertible, then

$$(A + B)^{-1} = A^{-1} - A^{-1}B(A + B)^{-1}.$$

Observe now that the the second-order lattice Laplacian has the form

$$\Delta = \sum_{(x, x') : \|x - x'\| = 1} \Gamma_{x, x'}, \quad \Gamma_{x, x'} = |\delta_x\rangle\langle\delta_{x'}|,$$

so that $(\Gamma_{x, x'} f)(y) = \delta_{x, y} f(x')$. Given a box $\Lambda_\ell(u) \subset \Lambda$, the Laplacian Δ_Λ in Λ with Dirichlet boundary conditions on $\partial^+ \Lambda$ reads as follows:

$$\Delta_\Lambda = \left(\Delta_{\Lambda_\ell(u)} \oplus \Delta_{\Lambda_\ell^c(u)} \right) + \sum_{(x, x') \in \partial \Lambda_\ell(u)} (\Gamma_{x, x'} + \Gamma_{x', x}).$$

Similarly, for the Hamiltonian H_Λ (with Dirichlet boundary conditions) we can write:

$$H_\Lambda = \left[(\Delta_{\Lambda_\ell(u)} + V_{\Lambda_\ell(u)}) \oplus (\Delta_{\Lambda_\ell^c(u)} + V_{\Lambda_\ell^c(u)}) \right] + \sum_{(x, x') \in \partial \Lambda_\ell(u)} (\Gamma_{x, x'} + \Gamma_{x', x}),$$

with $\Lambda_\ell^c(u) = \Lambda_L(u) \setminus \Lambda_\ell(u)$. Therefore, by the resolvent identity combined with the above decomposition, we have:

$$G(u, y; E) = \sum_{(x, x') \in \partial \Lambda_\ell(u)} G(u, x; E) \Gamma_{x, x'} G(x', y; E),$$

usually called the Geometric Resolvent Identity, yielding immediately the Geometric Resolvent Inequality (referred to as GRI in what follows):

$$|G(u, y; E)| \leq \left(\max_{x \in \partial^- \Lambda_\ell(u)} |G(u, x; E)| \cdot |\partial^+ \Lambda_\ell(u)| \right) \cdot \max_{x' \in \partial^+ \Lambda_\ell(u)} |G(x', y; E)|. \quad (2.5)$$

Throughout this paper, we use a standard notation $[[a, b]] := [a, b] \cap \mathbb{Z}$.

3. Green functions in a finite volume

Definition 3.1. A box $\Lambda_L(u)$ is called (E, m) -non-singular $((E, m)$ -NS) if

$$\max_{y \in \partial \Lambda_L(u)} |G(u, y; E)| \leq e^{-\gamma(m, L)},$$

where

$$\gamma(m, L) = m(L + L^{3/4}) = m \left(1 + L^{-1/4}\right) L. \quad (3.1)$$

Otherwise, it is called (E, m) -singular $((E, m)$ -S).

Remark. The function $\gamma(m, L)$ defined in Eqn (3.1) will be often used below. It allows us to avoid a "massive rescaling of the mass", which would, otherwise, inevitably make notations and assertions more cumbersome. Obviously, for large values of L , $\gamma(m, L)/(mL) \approx 1$.

It is convenient to introduce the following property (or assertion), the validity of which depends upon parameters $u \in \mathbb{Z}^d, L \in \mathbb{N}^*, m > 0, p > 0$, as well as upon the probability distribution of the random potential $\{V(x; \omega), x \in \mathbb{Z}^d\}$:

$$(\mathbf{S}, \mathbf{L}, \mathbf{m}) \quad \mathbb{P} \{ \Lambda_L(u) \text{ is } (E, m)\text{-S} \} \leq L^{-p}.$$

In order to distinguish between single- and multi-particle models, we will often write $(\mathbf{S}, \mathbf{L}, \mathbf{k}, \mathbf{N})$ where $N \geq 1$ is the number of particles.

Definition 3.2. A lattice subset $A \subset \mathbb{Z}^d$ of diameter L is called E -non-resonant (E -NR) if $\|G_A(E)\| \leq e^{-L^\beta}$, $\beta \in (0, 1)$, and E -resonant (E -R), otherwise. It is called (E, ℓ) -completely non-resonant $((E, \ell)$ -CNR) if it is E -NR and does not contain any E -R cube A' of diameter $\geq 3\ell$.

Lemma 3.1. If the marginal CDF $F(s) = F_V(s)$ is Hölder-continuous, then

$$\exists L^* > 0, \beta' \in (0, 1) : \forall L \geq L^* \mathbb{P} \left\{ \Lambda_L(u) \text{ is not } (E, L^{2/3})\text{-CNR} \right\} \leq e^{-L^{\beta'}}.$$

Consider a pair of boxes $\Lambda_\ell(u) \subset \Lambda_L(x)$. If $\Lambda_\ell(u)$ is (E, m) -NS, then GRI implies that function $f(x) = f_y(x) := G(x, y; E)$ satisfies

$$|f(u)| \leq \tilde{q} \cdot \max_{v: \|v-u\|=\ell} |f(v)|, \quad (3.2)$$

with

$$\tilde{q} = \tilde{q}(d, \ell; E) = 2d\ell^{d-1}e^{-\gamma(m, \ell)}. \quad (3.3)$$

Now suppose that $\Lambda_\ell(u)$ is (E, m) -S, but $\Lambda_L(x)$ is E -CNR and, in addition, for some $A > 0$ and for any w with $\text{dist}(w, \Lambda_{A\ell}(u)) = \ell$ the box $\Lambda_\ell(w)$ is (E, m) -NS. Then, by GRI applied twice,

$$\begin{aligned} |G(u, y)| &\leq 2d(6\ell)^{d-1}e^{L^\beta} \max_{w: \|w-u\|=(A+1)\ell} |G(w, y; E)| \\ &\leq 2d(6\ell)^{d-1}e^{L^\beta} 2d(2\ell)^{d-1}e^{-\gamma(m, \ell)} \max_{v: \|v-u\| \in [A\ell, (A+2)\ell]} |G(v, y; E)|. \end{aligned}$$

Therefore, with

$$q(d, \ell, E) := 4d^2(12\ell^2)^{d-1}e^{L^\beta}e^{-\gamma(m, \ell)} > \tilde{q}(d, \ell, E), \quad (3.4)$$

we obtain

$$|G(u, y)| \leq \tilde{q}(d, \ell, E) \max_{v: \|v-u\| \in [A\ell, (A+2)\ell]} |G(v, y; E)|. \quad (3.5)$$

With these observations in mind, we study in the next section decay properties of functions $f : \Lambda_L(x) \rightarrow \mathbb{C}$ obeying, for any point $u \in \Lambda_L(x)$, one of the Eqns (3.2), (3.5). Observe that we do not require that q and $\tilde{q}$ be smaller than 1, but, of course, the above bounds (and those obtained in Section 4) are useful only for $q, \tilde{q} < 1$. Finally, note that, since $q < \tilde{q}$, it will be convenient to replace $\tilde{q}$ by q in the bound (3.2). With this modification, we see that the only difference between bounds (3.2) and (3.5) is in the distance $\|v - u\|$ figuring in these inequalities.

4. Radial descent: A few simple lemmas

Definition 4.1. Consider a set $\Lambda \subset \mathbb{Z}^d$ (not necessarily finite), a subset $\mathcal{S} \subset \Lambda$ and a bounded function $f : \Lambda \rightarrow \mathbb{C}$. Let $L \geq 0$ be an integer and $q > 0$. Function f will be called (ℓ, q) -subharmonic in Λ and regular on $\Lambda \setminus \mathcal{S}$ (or, equivalently, $(\ell, q, \mathcal{S})$ -subharmonic), if for any $u \notin \mathcal{S}$ with $\text{dist}(u, \partial\Lambda) \geq \ell$, we have

$$|f(u)| \leq q \max_{y: \|y-u\|=\ell} |f(y)| \quad (4.1)$$

and for any $u \in \mathcal{S}$

$$|f(u)| \leq q \max_{y: \text{dist}(y, \mathcal{S}) \in [1, 2\ell-1]} |f(y)|. \quad (4.2)$$

On Fig.1, the pink square (labeled by the letter $\mathcal{S}$) is a singular set, and the green belt-shaped area is formed by non-singular (NS) neighboring squares (the four white squares are examples of neighboring NS-squares). A thin white layer between the pink singular square and its NS-neighborhood has the width = 1 (the minimal distance on the lattice between two disjoint sets).

For our purposes, it suffices to consider a particular case where $\Lambda = \Lambda_L(x)$ is a cube of side L with center x and $\mathcal{S} = \Lambda_{\tilde{\ell}}(v) \cap \Lambda$, $\tilde{\ell} = A\ell - 1$. In this particular case Eqn (4.2) becomes

$$\forall u \in \mathcal{S} \quad |f(u)| \leq q \max_{y: \ell \leq \|y-u\| \leq A\ell-1} |f(y)|. \quad (4.2A)$$

We will use the notation $\mathcal{M}(f, \Lambda) := \max_{x \in \Lambda} |f(x)|$. Our goal is to obtain an upper bound on the value $f(x)$ exponential in L/ℓ . To this end, we study separately several cases.

Lemma 4.1. Let $\Lambda = \Lambda_L(x)$ and consider an $(\ell, q, \emptyset)$ -subharmonic function $f : \Lambda \rightarrow \mathbb{C}$. Then

$$|f(x)| \leq q^{\lfloor L/\ell \rfloor - 1} \mathcal{M}(f, \Lambda). \quad (4.3)$$

Proof. Let $n \geq 2$ and consider a point $u \in \Lambda_{L-n\ell}(x)$. Since $\Lambda_\ell(u) \subset \Lambda_{L-(n-1)\ell}(x)$, the subharmonicity condition implies that

$$|f(u)| \leq q \max_{y \in \Lambda_L(x): \|y-u\|=\ell} |f(y)| \leq q \mathcal{M}(f, \Lambda_{L-(n-1)\ell}(x)). \quad (4.4)$$

In other words, we have

$$\mathcal{M}(f, \Lambda_{L-n\ell}(x)) \leq q \mathcal{M}(f, \Lambda_{L-(n-1)\ell}(x)). \quad (4.5)$$




Fig. 1. An example of a singular subset (pink). All squares inside the green area are non-singular. Four white squares are examples of non-singular squares

Eqn (4.5) can be iterated, as long as the set $\Lambda_{L-n\ell}(x)$ is non-empty, so we obtain by induction

$$\mathcal{M}(f, \Lambda_{L-n\ell}(x)) \leq q^{n-1} \mathcal{M}(f, \Lambda_{L-\ell}(x)) \leq q^{n-1} \mathcal{M}(f, \Lambda_L(x)). \quad (4.6)$$

Now the assertion of the lemma follows from the inclusion $x \in \Lambda_{L-[L/\ell]\ell}(x)$. $\square$

Lemma 4.2. *Let $\Lambda = \Lambda_L(x)$ and consider an $(\ell, q, \mathcal{S})$ -subharmonic function $f : \Lambda \rightarrow \mathbb{C}$. Suppose that $\text{diam}(\mathcal{S}) \leq A\ell - 1$ and $\text{dist}(\mathcal{S}, \partial\Lambda_L(x)) < \ell$. Then*

$$|f(x)| \leq q^{[(L-A)/\ell]-2} \mathcal{M}(f, \Lambda). \quad (4.7)$$

Proof. Observe that $\mathcal{S} \subset (\Lambda_L(x) \setminus \Lambda_{L-(A+1)\ell}(x))$. Therefore, function f is $(L, q, \emptyset)$ -subharmonic in $\Lambda_{L-(A+1)\ell}(x)$, and it suffices to apply Lemma 4.1. $\square$

Lemma 4.3. *Let f be an $(\ell, q, \mathcal{S})$ -subharmonic function on $\Lambda = \Lambda_L(x)$. Suppose that $\text{diam}(\mathcal{S}) = A\ell - 1$ and*

$$(r+2)\ell \leq \text{dist}(\mathcal{S}, \partial\Lambda_L(x)) < (r+3)\ell,$$

so that

$$\mathcal{S} \subset \Lambda_{L-(r+2)\ell}(x) \setminus \Lambda_{L-(r+A+3)\ell}(x).$$

Then

$$|f(x)| \leq q^{[(L-A)/\ell]-3} \mathcal{M}(f, \Lambda). \quad (4.8)$$




Fig. 2. "Radial" induction in Lemma 4.2. Here, a singular subset (pink) is at distance $< \ell$ from the boundary, and it suffices to start the induction from the inner square (its boundary is indicated by the double line). The singular set can be here even an "incomplete" square.

Proof. We start as in Lemma 4.1. For points $u \in \Lambda_{L-\ell}(x) \setminus \mathcal{S}$ we use Eqn (4.1):

$$|f(u)| \leq q \max_{y: |y-u|=L} |f(y)| \leq q \mathcal{M}(f, \Lambda_L(x)),$$

and for points $u \in \mathcal{S} \subset \Lambda_{L-\ell}(x)$ Eqn (4.2) leads to a similar upper bound:

$$|f(u)| \leq q \max_{y: |y-u| \in [A\ell, (A+2)\ell-1]} |f(y)| \leq q \mathcal{M}(f, \Lambda_L(x)).$$

Therefore,

$$\mathcal{M}(f, \Lambda_{L-\ell}(x)) \leq q \mathcal{M}(f, \Lambda_L(x)). \quad (4.9)$$

We can iterate this argument and obtain, for $n = 1, \dots, r$

$$\mathcal{M}(f, \Lambda_{L-n\ell}(x)) \leq q \mathcal{M}(f, \Lambda_{L-(n-1)\ell}(x)). \quad (4.10)$$

Indeed, if $1 \leq n \leq r$, then

$$\{u : \text{dist}(\mathcal{S}, u) \in [1, 2\ell - 1]\} \subset \Lambda_{L-n\ell}(x) \subseteq \Lambda_{L-r\ell}(x).$$

This leads to the following upper bound:

$$\mathcal{M}(f, \Lambda_{L-r\ell}(x)) \leq q^r \mathcal{M}(f, \Lambda_L(x)). \quad (4.11)$$




Fig. 3. Two-fold "radial" induction in Lemma 4.3. Here, a singular subset (pink) is at distance $r' \in [(r+2)\ell, (r+3)\ell]$ from the boundary.

If $L - (A + r + 3)\ell < 2\ell$, we stop the induction and obtain for $x \in \Lambda(x)$ the required upper bound.

Otherwise, we make at least one inductive step (or more) inside the box $\Lambda_{L-(r+A+3)\ell}(x)$. Since $\Lambda_{L-(r+A+3)\ell}(x) \subset (\Lambda_{L-r\ell}(x) \setminus \mathcal{S})$, Eqn (4.11) implies

$$\mathcal{M}(f, \Lambda_{L-(r+A+3)\ell}(x)) \leq q^r \mathcal{M}(f, \Lambda_L(x)). \quad (4.12)$$

Observe that function f is $(\ell, q, \emptyset)$ -subharmonic on $\Lambda_{L-(r+A+3)\ell}(x)$. Applying Lemma 4.1 to this cube, we conclude that

$$|f(x)| \leq q^{[(L-r-A)/\ell]-3} q^r \mathcal{M}(f, \Lambda_L(x)) \leq q^{[(L-A)/\ell]-3} \mathcal{M}(f, \Lambda_L(x)). \quad \square$$

5. Inductive bounds of Green functions

Taking into account observations made at the end of Section 3, namely, Eqns (3.2) and (3.5), we come immediately to the following

Lemma 5.1. *Suppose that a box $\Lambda_L(u)$ is E -NR and does not contain any (E, m) -S box of size ℓ . Then, with $q = q(d, \ell, E)$ defined in Eqn (3.12),*

$$\max_{y \in \partial^- \Lambda_L(u)} |G_\Lambda(u, y; E)| \leq q^{[L/\ell]} \|G_\Lambda(E)\|. \quad (5.2)$$

In particular, if ℓ^{-1} and ℓ/L are sufficiently small, then $\forall y \in \partial A_L(u)$, we have

$$|G_A(u, y; E)| \leq e^{-\gamma(m, L)} \|G_A(E)\|. \quad (5.3)$$

Proof. Apply Lemma 4.3. $\square$

Lemma 5.2. *Suppose that a box $A_L(u)$ is E -CNR and does not contain any pair of non-overlapping (E, m) -S boxes of size ℓ . Then*

$$\max_{y \in \partial A_L(u)} |G_{A_L}(u, y)| \leq e^{-\gamma(m, L)}. \quad (5.4)$$

Proof. If $A_L(u)$ contains no (E, m) -S box $A_\ell(x)$, the assertion follows from Lemma 5.1. Suppose there exists a box $A_\ell(x) \subset A_L(u)$ which is (E, m) -S. Since all boxes $A_\ell(w)$ with $\|w - x\| = 2\ell$ are disjoint with $A_\ell(x)$, none of them is (E, m) -S, by hypotheses of the lemma. Therefore, either Lemma 4.2 or Lemma 4.3 applies, with $A = 4$, giving the required upper bound. $\square$

Lemma 5.3. *Suppose that the finite-volume approximations $H_A(\omega)$, $A \subset \mathbb{Z}^d$, $|A| < \infty$, of random LSO $H(\omega)$ in $\ell^2(\mathbb{Z}^d)$ satisfy the following hypotheses:*

(W) $\mathbb{P} \left\{ \text{dist}[E, \Sigma(H_A(\omega))] \leq e^{-|A|^{-\beta}} \right\} \leq e^{-|A|^{-\beta'}}$, for some $\beta, \beta' > 0$;

(DS. ℓ) $\forall v$ such that $A_\ell(v) \subset A_L(u)$, $\mathbb{P} \{ A_\ell(u) \text{ is } (E, m)\text{-S} \} \leq \ell^{-p}$.

Set $L = \lceil \ell^{3/2} \rceil$. If $p > 6d$, then H_{A_L} satisfies **(S.L, $m, 1$)**.

Proof. Consider a box $A_L(u)$. By Lemma 5.2, it must be (E, m) -NS, unless one of the following events occurs:

- $A_L(u)$ is not E -CNR
- $A_L(u)$ contains at least two non-overlapping (E, m_k) -S boxes $A_\ell(x), A_\ell(y)$.

The probability of the former event is bounded by **(W)**. Further, by virtue of **(DS. ℓ)** the probability of the latter event is bounded by

$$\frac{1}{2} L^{2d} \ell^{-2p} \leq \frac{1}{2} L^{-p \left(\frac{2}{372} - \frac{2d}{p} \right)} \leq L^{-p}. \quad \square$$

Corollary 5.1. **(S.L $_k, m, 1$)** implies **(S.L $_{k+1}, m, 1$)** (provided that Wegner-type bound **(W)** holds true).

Theorem 5.1. *If an IID random potential $V(x; \omega)$ satisfies the assumption (2.2), then **(S.L $_k, 1$)** holds true for all $k \geq 0$.*

In the single-particle case, treated in this section, for Hamiltonians with an IID potential, it is well-known that the exponential decay of Green functions implies that the respective Hamiltonian has pure point spectrum, and that all its eigenfunctions decay exponentially (with probability one). This follows from Simon–Wolff criterion (cf. [SW86]), which applies also to a large class of so-called non-deterministic potentials. However, the method of [SW86] does apply to multi-particle Hamiltonians. For this reason, the results of the next section do not lead directly to spectral localization for multi-particle Hamiltonians.

6. Simplified multi-particle MSA

In this section, we study decay properties of Green functions for an N -particle Hamiltonian $H^{(N)}(\omega)$ defined in Eqn (2.4). We stress that estimates given below are *far from optimal*; they only show that for any given number of particles $N \geq 1$ and any $m > 0$, there exists a threshold $g_N = g_N(m) > 0$ for the disorder parameter g such that if $|g| \geq g_N$, then Green functions in the N -particle model (with a short-range interaction) decay exponentially with rate m . It is also important to realize that using the MSA, in its traditional form, for an N -particle system with large N inevitably requires using large values of g_N . Indeed, this phenomenon occurs even in the single-particle MSA in high dimension $d \gg 1$: the respective threshold $g_1 = g_1(m, d) \rightarrow \infty$ as $d \rightarrow \infty$. A direct inspection of the conventional MSA show that, typically, $V(\cdot; \omega, g_1(m, d)) = O(d)$.

Given a number $m > 0$ and an integer $L_0 > 0$, we will define a decreasing sequence of decay exponents $m^{(n)}$, $n \geq 1$ as follows:

$$m^{(1)} = m; \quad m^{(N)} = m^{(N-1)} - L_0^{-1/8}, \quad N > 1.$$

It is clear that in order to have $m^{(N)} > 0$, we have to assume that $m = m^{(1)}$ is sufficiently large (depending on N). In turn, this requires the thresholds $g_1, \dots, g_{N-1}$ to be large enough, as explained above.

Compared to the single-particle MSA scheme presented in previous sections, we have to replace the property **(S.L, I)** by a different one, **(MS.L, I, N)** given below, and to include in **(MS.L, I, N)** the requirement that the decay of Green functions holds for all $n' < N$:

(MS.L_k, I, N): The property **(S.L_k, I, n)** holds for $1 \leq n \leq N - 1$,

$$\mathbb{P} \left\{ \Lambda_L^{(n)}(\mathbf{u}) \text{ is } (E, m^{(n)})\text{-S} \right\} \leq L^{-p(n, g)},$$

where $p(n, g) \rightarrow \infty$ as $|g| \rightarrow \infty$, $n = 1, \dots, N - 1$, and $E \in I$. In the case of large disorder, one can set $I = \mathbb{R}$, and in the case of "low energies", I is a sufficiently small interval near the bottom of the spectrum.

Remark. For any $k = 0$ and any $n > 1$, the validity of the above statement is proved exactly in the same way as for $n = 1$. Then for $k > 0$ it is reproduced inductively. Denoting $P(N - 1, g) = \min_{1 \leq n \leq N-1} p(n, g)$, we see that, under the hypothesis **(MS.L_k, I, N)**, we have $P(N - 1, g) \rightarrow \infty$ as $|g| \rightarrow \infty$.

For the sake of notational simplicity, we consider in Lemma 6.1 below only partitions $[[1, N]] = \mathcal{J} \amalg \mathcal{J}^c$ of the interval $[[1, N]]$ into two (non-empty) consecutive sub-intervals, $\mathcal{J} = [[1, n']]$, $\mathcal{J}^c = [[n' + 1, N]]$. In other words, we consider a union of two subsystems with particles $1, \dots, n'$ and $n' + 1, \dots, N$, respectively.

We introduce the "diagonal" subset of $\mathbb{Z}^{Nd}$

$$\mathbb{D} := \left\{ x = (x_1, \dots, x_1), x_1 \in \mathbb{Z}^d \right\}.$$

Lemma 6.1. *Let $[[1, N]] = \mathcal{J} \cup \mathcal{J}^c$, $\mathcal{J} = [[1, n']]$, $\mathcal{J}^c = [[n' + 1, N]]$ and consider a cube $\Lambda_L^{(N)}(\mathbf{u}) = \Lambda_L^{(n')}(u') \times \Lambda_L^{(n'')}(u'')$, with $u' = (u_1, \dots, u_{n'})$, $u'' = (u_{n'+1}, \dots, u_N)$. Let $\Sigma' = \{\lambda_a, a = 1, \dots, |\Lambda^{(n')}(u')|\}$ be the spectrum of $H_{\Lambda^{(n')}(u')}^{(n')}$, and $\Sigma'' = \{\mu_b, b = 1, \dots, |\Lambda^{(n'')}(u'')|\}$ the spectrum of $H_{\Lambda^{(n'')}(u'')}^{(n'')}$. Suppose that*

(i) *the interaction between these two subsystems vanishes:*

$$\forall j \in \mathcal{J}, \forall i \in \mathcal{J}^c \quad \|u_j - u_i\| > 2(L-1) + r_0, \quad (6.1A)$$

so that

$$\forall v_j \in \Lambda_L^{(1)}(u_j), \forall v_i \in \Lambda_L^{(1)}(u_i) \quad U_2(v_j, v_i) = 0. \quad (6.1B)$$

(ii) $\Lambda_L^{(N)}(\mathbf{u})$ is E -NR;

(iii) for any $\mu_b \in \Sigma'$, $\Lambda_L^{(n')}(u')$ is $(E - \mu_b, m^{(N-1)})$ -NS, and for any $\lambda_a \in \Sigma'$, $\Lambda_L^{(n'')}(u'')$ is $(E - \lambda_a, m^{(N-1)})$ -NS.

Then $\Lambda_L^{(N)}(\mathbf{u})$ is $(E, m^{(N)})$ -NS.

The proof of Lemma 6.1 is given in Section 7; it is fairly straightforward.

We will call N -particle boxes $\Lambda_L^{(N)}(u)$ admitting a decomposition described in Lemma 6.1 *decomposable*. It is easy to see that an N -particle box $\Lambda_L^{(N)}(u)$ is non-decomposable if and only if the union of cubes

$$\bigcup_{j=1}^N \Lambda_{L+r_0/2}^{(1)}(u_j) \subset \mathbb{R}^{Nd}$$

is connected (for this argument, we identify $\Lambda_L^{(1)}(u_j)$ with a cube in $\mathbb{R}^d$ of side $2(\ell-1)$ with center u_j , by a slight abuse of notations). In turn, such a union is connected, then

$$\forall j = 2, \dots, N \quad \|u_j - u_1\| \leq 2(L-1) + r_0,$$

so that

$$\text{dist}(u, \mathbb{D}) \leq \|u - (u_1, \dots, u_1)\| \leq 2N(L-1) + Nr_0 =: r_{N,L}.$$

Clearly, if $\text{dist}(u, \mathbb{D}) > r_{N,L}$, then $\Lambda_L^{(N)}(u)$ is decomposable. Observe that in this case the families of random variables

$$\mathcal{V}' := \{V(x; \omega), x \in \bigcup_{j=1}^{n'} \Lambda_L^{(1)}(u_j)\}, \quad \mathcal{V}'' := \{V(x; \omega), x \in \bigcup_{j=n'+1}^N \Lambda_L^{(1)}(u_j)\}$$

are mutually independent, and so are the random spectra $\Sigma' = \Sigma'(\omega)$, $\Sigma'' = \Sigma''(\omega)$. Also, observe that if EVs $\{\lambda_a\}$ and $\{\mu_b\}$ are non-negative, then $E - \lambda_a \leq E$ and $E - \mu_b \leq E$. Therefore, in a situation where $I = [E_0^*, E_0^* + E_1^*]$ is a bounded interval, and we attempt to prove the decay of Green functions only for "sufficiently low energies", i.e. for E_1^* small enough, then the hypotheses on Green functions for subsystems with smaller number of particles ($n', n'' < N$) required for the induction on the particle number ($N-1 \rightsquigarrow N$) are formulated for energies E which either belong to $[E_0^*, E_0^* + E_1^*]$ or are beyond the spectra of Hamiltonians $H_{\Lambda^{(n')}(u')}$ and/or $H_{\Lambda^{(n'')}(u'')}$. More precisely, in the latter case E is *below* the ground state energy. This allows using the well-known Combes-Thomas argument in order to prove the exponential decay of Green functions for Hamiltonians $H_{\Lambda^{(n')}(u')}$ and/or $H_{\Lambda^{(n'')}(u'')}$. In the case of a strong disorder, one can simply take $I = \mathbb{R}$ for Hamiltonians $H_{\Lambda^{(n')}(u')}$ and $H_{\Lambda^{(n'')}(u'')}$.

Lemma 6.2. *Fix an integer $N > 1$ and suppose that $(\mathbf{MS.L}_k, \mathbf{I}, \mathbf{n})$ holds for all $n < N$. If an N -particle box $\Lambda^{(N)}(u)$ is decomposable, then*

$$\mathbb{P} \left\{ \Lambda_{L_k}^{(N)}(u) \text{ is } (E, m)\text{-S} \right\} \leq L_k^{-P(N-1, g)} + e^{-L_k^\beta}.$$

Therefore, for any $P > 0$ and for all $|g|$, L_0 large enough, we have

$$\mathbb{P} \left\{ \Lambda_{L_k}^{(N)}(u) \text{ is } (E, m)\text{-S} \right\} \leq L_k^{-P}. \quad (4.3)$$

Proof. Since $\Lambda_{L_k}^{(N)}(\mathbf{u})$ is decomposable, operator $H_{\Lambda_{L_k}^{(N)}(\mathbf{u})}^{(N)}$ admits representation (7.1) for some $n', n'' < N$. By Lemma 6.1, either

- the box $\Lambda_{L_k}^{(N)}(\mathbf{u})$ is E -R, which occurs with probability $\leq e^{-L_k^\beta}$, or
- the box $\Lambda_{L_k}^{(n')}(u')$ is $(E - \mu_b, m_k)$ -S, for some EV μ_b of operator $H_{\Lambda_{L_k}^{(n'')}(u'')}^{(n'')}$, or
- the box $\Lambda_{L_k}^{(n')}(u')$ is $(E - \lambda_a, m_k)$ -S, for some EV λ_a of operator $H_{\Lambda_{L_k}^{(n')}(u')}^{(n')}$.

The latter two events have probabilities $\leq L_k^{-P(N-1, g) + N - 1}$, by $(\mathbf{MS.L}_k, \mathbf{I}, N - 1)$. Indeed, since the projection boxes $\Lambda_{L_k}^{(n')}(u')$ and $\Lambda_{L_k}^{(n'')}(u'')$ are disjoint, we can condition on one of them, and estimate the conditional probability, where a fixed-energy argument applies. $\square$

Now we are prepared to give a proof of the main result of the fixed-energy MSA. For notational simplicity, below we write m instead of $m^{(N)}$.

Theorem 6.1. *Fix an interval $I \subset \mathbb{R}$ and an integer $N > 1$. Then, for $P(N - 1, g)$ large enough, the property $(\mathbf{MS.L}_k, \mathbf{I}, N)$ implies $(\mathbf{MS.L}_{k+1}, \mathbf{I}, N)$.*

Proof. By Wegner bound, $\mathbb{P} \left\{ \Lambda_{L_{k+1}}^{(N)} \text{ is not } E\text{-CNR} \right\} \leq e^{-L_{k+1}^\beta}$. Further, set

$$\mathbf{S} = \left\{ \exists \Lambda_{L_k}(x) \subset \Lambda_{L_{k+1}}(u) \text{ which is } (E, m)\text{-S and } \text{dist}(\Lambda_{L_k}(x), \mathbb{D}) \leq r_{N, L} \right\}.$$

By Lemma 6.2, and for $P(N - 1, g)$ large enough, $\mathbb{P} \{ \mathbf{S} \} \leq \frac{1}{2} L_{k+1}^{-p(N, g)}$. Next, set

$$\mathbf{B} = \left\{ \exists \Lambda_{L_k}(x), \Lambda_{L_k}(y) \subset \Lambda_{L_{k+1}}(u) \text{ which are } (E, m)\text{-S and such that } \|x - y\| > 9(L_k + r_{N, L_k}), \text{dist}(\Lambda_{L_k}(x), \mathbb{D}) \leq r_{N, L}, \text{dist}(\Lambda_{L_k}(y), \mathbb{D}) \leq r_{N, L} \right\}.$$

Fix centers x, y . By Lemma 8.1, potential samples in boxes $\Lambda_{L_k}(x), \Lambda_{L_k}(y)$ close to the "diagonal" $\mathbb{D}$ are **independent**, so that

$$\mathbb{P} \{ \mathbf{B} \} \leq \frac{1}{2} L_{k+1}^{2d} \left(\max_{v \in \Lambda_{L_{k+1}}(u)} \mathbb{P} \{ \Lambda_{L_k}(v) \text{ is } (E, m)\text{-S} \} \right)^2 < \frac{1}{2} L_{k+1}^{-p(N, g)}.$$

If neither of events $\mathbf{S}, \mathbf{B}$ occurs, then, by Lemma 5.3, $\Lambda_{L_{k+1}}(\mathbf{u})$ is (E, m_{k+1}) -NS. Therefore, with $p(N, g) \rightarrow \infty$ as $|g| \rightarrow \infty$, we have

$$\mathbb{P} \left\{ \Lambda_{L_{k+1}}^{(N)}(\mathbf{u}) \text{ is } (E, m)\text{-S} \right\} \leq \frac{1}{2} L_{k+1}^{-p(N, g)} + \frac{1}{2} L_{k+1}^{-p(N, g)} = L_{k+1}^{-p(N, g)}. \quad \square$$

7. Appendix A

Proof of Lemma 6.1

Let $\{\varphi_a, \lambda_a\}$ be the normalized eigenfunctions (EFs) and the respective eigenvalues (EVs) of $H_{\Lambda^{(n')}(u')}$, and $\{\psi_b, \mu_b\}$ be the normalized EFs and the EVs of $H_{\Lambda^{(n'')}(u'')}$. Due to absence of interaction between configurations u', u'' , we have

$$H_{\Lambda^{(N)}(\mathbf{u})}^{(N)} = H_{\Lambda^{(n')}(u')}^{(n')} \otimes \mathbf{1} + \mathbf{1} \otimes H_{\Lambda^{(n'')}(u'')}^{(n'')}. \quad (7.1)$$

Therefore, the EFs $\Psi_{a,b}$ of operator $H_{\Lambda^{(N)}(\mathbf{u})}^{(N)}$ have the form $\Psi_{a,b} = \varphi_a \otimes \psi_b$, and the respective EVs are given by $E_{a,b} = \lambda_a + \mu_b$, so that $E - E_{a,b} = (E - \lambda_a) - \mu_b$.

Consider any point $\mathbf{v} \in \partial \Lambda_L^{(N)}(\mathbf{u})$. Then either $\|u' - v'\| = L$, or $\|u'' - v''\| = L$. W.l.o.g., assume that $\|u' - v'\| = L$. Since $\|\varphi_a\| = 1$ and $n' < N$, we can write

$$\begin{aligned} |\mathbf{G}(\mathbf{u}, \mathbf{v}; E)| &\leq \sum_a |\varphi_a(u') \varphi_a(v')| \left| \sum_b \frac{\psi_b(u'') \psi_b(v'')}{(E - \lambda_a) - \mu_b} \right| \\ &\leq \left| \Lambda_L^{(n')}(u') \right| \max_{\lambda_a} |G_{\Lambda_L^{(n'')}(u'')}(u'', v''; E - \lambda_a)| \leq L^{N-1} e^{-L^\beta} e^{-mL}. \quad \square \end{aligned}$$

8. Appendix B

Lemma 8.1. *Consider two points $x, y \in \mathbb{Z}^{Nd}$ and an integer $L > 0$. Suppose that $\|x - y\| > AL$ with $A = 4B + 6$, $B > 0$, and*

$$\text{dist}(\Lambda_L^{(N)}(x), \mathbb{D}_0) < BL \quad \text{dist}(\Lambda_L^{(N)}(y), \mathbb{D}_0) < BL. \quad (8.1)$$

Then for all $j = 1, \dots, N$, we have $\Pi_j \Lambda_L^{(N)}(x) \cap \Pi_j \Lambda_L^{(N)}(y) = \emptyset$.

Proof. (See Fig. 4 illustrating the case $N = 2, d = 1$.) Eqn (8.1) implies that there exist some points $u = (u_1, \dots, u_1) \in \mathbb{D}$, $v = (v_1, \dots, v_1) \in \mathbb{D}$ such that $\|x - u\| \leq L + BL$, $\|y - v\| \leq L + BL$. Then we have, by triangle inequality,

$$\|x - y\| \leq \|x - u\| + \|u - v\| + \|v - y\| \leq (2B + 2)L + \|u - v\|,$$

so that

$$\|u - v\| \geq \|x - y\| - (2B + 2)L > 2(B + 2)L, \quad (8.2)$$

and also

$$\Lambda_L(x) \subset \Lambda_{(B+2)L}(u), \quad \Lambda_L(y) \subset \Lambda_{(B+2)L}(v). \quad (8.3)$$

Similar inclusions hold true for the projections:

$$\Pi_j \Lambda_L^{(N)}(x) \subset \Pi_j \Lambda_{(B+2)L}(u), \quad \Pi_j \Lambda_L^{(N)}(y) \subset \Pi_j \Lambda_{(B+2)L}(v).$$

However, Eqn (8.2) implies that cubes $\Lambda_{(B+2)L}(u)$ and $\Lambda_{(B+2)L}(v)$ are disjoint, and so are, therefore, the projections $\Pi_j \Lambda_L^{(N)}(x)$ and $\Pi_j \Lambda_L^{(N)}(y)$. $\square$

Acknowledgments.

I thank the Isaac Newton Institute (INI) and the organizers of the INI programs IGCS (2003), AGA (2007) and MPA (2008) for their kind invitations




Fig. 4. Cubes $\Lambda_L^{(2)}(x)$ and $\Lambda_L^{(2)}(y)$ with non-overlapping projections

and warm hospitality during my stay at the INI. It is my pleasure to thank also many participants to these programs, especially MPA, for numerous fruitful discussions. In fact, the main idea of this short note is a direct result of my discussions with Boris Shapiro. I regret that Boris did not find possible to become a co-author of the present paper; it does not make, however, his contribution less important. Special thanks to Tom Spencer for initiating my collaboration with Boris Shapiro and many stimulating discussions, and also to Y. Fyodorov, A. Mirlin, D. Shepelyanski and S. Fishman (in chronological order of discussions) for sharing with me their physical insights.

I thank Yuri Suhov, with whom we have been running since five years, by Yuri's initiative, a larger project aimed to establish the Anderson localization phenomenon for multi-particle systems, for stimulating discussions.

I also thank Anne Boutet de Monvel and Peter Stollmann for discussions of a preliminary version of this short note and making a number of useful suggestions.

References

- [AW08] M. Aizenman, S. Warzel, *Localization Bounds for Multiparticle Systems*. - Preprint, arXiv math-ph/0809:3436, 2008.
- [BCSS08] A. Boutet de Monvel, V. Chulaevsky, P. Stollmann, Y. Suhov: *Wegner-type bounds for a two-particle continuous Anderson model with an alloy-type external potential*. - arXiv:math-ph/0821:2621, 2008.

- [BCS08] A. Boutet de Monvel, V. Chulaevsky, Y. Suhov: *Wegner-type bounds for a two-particle Anderson model in a continuous space*. - arXiv:math-ph/0821:2627, 2008.
- [CS08] V. Chulaevsky, Y. Suhov: *Wegner bounds for a two-particle tight binding model*. - Commun. Math. Phys., **283**, 479–489, 2008.
- [CS09A] V. Chulaevsky, Y. Suhov: *Eigenfunctions in a two-particle Anderson tight binding model*. - To appear in: Commun. Math. Phys.
- [CS09B] V. Chulaevsky, Y. Suhov: *Multi-Particle Anderson Localisation: Induction on the Number of Particles*. - Preprint, arXiv math-ph/ 0811:2530, 2008.
- [Dr87] H. von Dreifus, *On the effect of randomness in ferromagnetic models and Schrödinger operators*. PhD thesis, New York University, 1987.
- [DK89] H. von Dreifus, A. Klein: *A new proof of Localization in the Anderson Tight Binding Model*. - Commun. Math. Phys., **124**, 285–299, 1989.
- [FS83] J. Fröhlich, T. Spencer: *Absence of diffusion in the Anderson tight binding model for large disorder or low energy*. - Commun. Math. Phys., **88**, 151–184, 1983.
- [FMSS85] J. Fröhlich, F. Martinelli, E. Scoppola, T. Spencer: *A constructive proof of localization in Anderson tight binding model*. - Comm. Math. Phys., **101**, 21–46, 1985.
- [S88] T. Spencer: *Localization for Random and Quasiperiodic Operators*. - J. Statist. Phys., **51**, No. 5/6, 1009–1019, 1988.
- [St01] P. Stollmann, *Caught by disorder*. - Birkhäuser, 2001.
- [SW86] B. Simon, T. Wolff: *Singular Continuous Spectrum under Rank One Perturbations and Localization for Random Hamiltonians*. - Commun. Pure App. Math., **39**, 75–90, 1986.